

COGNITIVE IMPAIRMENT FOLLOWING USE OF ANAESTHETIC AGENTS: A REVIEW OF THE LITERATURE, AND IMPLICATIONS FOR FUTURE PRACTICE

Arjun Nesaratnam¹, Nisha Nesaratnam² & Mark Agius³

¹Addenbrooke's Hospital, Cambridge, Cambridge, UK

²School of Clinical Medicine, University of Cambridge, Cambridge, UK

³Clare College, University of Cambridge, Cambridge, UK

SUMMARY

Emerging data suggest that both inhalational and intravenous anaesthetics are associated with cognitive decline, particularly in the elderly. Of particular concern, is the phenomenon known as post-operative cognitive decline (POCD), characterised by a transient impairment of memory, concentration, language comprehension and social integration. Implicated in the pathophysiology of POCD is the accumulation of A β peptide, and inhibition of cholinergic neurotransmission. Currently used anaesthetic agents differ in their propensity for causing POCD as a result of their differing pharmacological profiles. Coupled with increasing knowledge of patient susceptibility factors, this information allows for modification of clinical practice to minimise the long-term disability that may otherwise accompany POCD.

Key words: post-operative cognitive dysfunction – anaesthesia - neurotoxicity

* * * * *

INTRODUCTION

Advances in surgery and anaesthesia have contributed to the 60% increase in the number of operations performed in the NHS over the last ten years. Whilst mortality attributable to general anaesthesia remains low (Jenkins et al. 2003), attention has been directed towards prevention of anaesthesia-associated morbidity, including cardiac, renal, cerebral and pulmonary sequelae. Emerging data suggest that both inhalational and intravenous anaesthetics are associated with cognitive decline, particularly in the elderly (Bekker et al. 2003). Of particular concern, is the phenomenon known as post-operative cognitive decline (POCD), characterised by a transient (days to weeks) impairment of memory, concentration, language comprehension and social integration (Phillips-Bute et al. 2006). Distinct from post-operative delirium due to its non-fluctuating course, retained consciousness levels and increased mortality rate, POCD may be affecting 15-25% patients undergoing surgery (Bryson et al. 2006). Nevertheless, many studies of POCD to date have lacked the statistical power to disentangle POCD from age-related cognitive decline, and indeed, to disentangle the effects of surgical trauma from that of anaesthetic agents (Dabrowski et al. 2012). Given the widespread use of anaesthetic agents in our aging population, it is crucial to understand the pathophysiology of anaesthetic-induced POCD, with a view to minimise any possible neurotoxicity.

PATHOPHYSIOLOGY OF POST-OPERATIVE COGNITIVE DECLINE

Although its pathophysiology has not been fully elucidated, it is likely that anaesthetic-induced POCD

has a multifactorial aetiology, involving a combination of genetic susceptibility, pre- and peri-operative factors. That cerebral vascular disease, systemic hypertension, diabetes mellitus and age are risk factors for POCD (Wolman et al. 1999) lends credence to the hypothesis that POCD is related to vascular insufficiency. Animal models reveal, however, a predominant role for neuroinflammation in the aetiology of POCD, with Mandal et al. (2009) noting widespread inflammation and A β peptide oligomerisation induced by the anaesthetic agents, isoflurane and desflurane. The increased incidence of POCD following surgical trauma, hypothermia and cardio-pulmonary bypass (Diegeler et al. 2000), may thus be a reflection of an increased peri-operative neuroinflammatory response.

The accumulation of A β peptide, loss of cholinergic forebrain neurons and positive predictive value of the apolipoprotein E-4 allele in POCD have naturally prompted comparisons with Alzheimer's disease (Xie et al. 2006). In both conditions, the impairments in cognition are likely related to a neurotoxic effect of amyloid β peptide, whose accumulation is increased following exposure to certain anaesthetic agents. Interestingly, Hellstrom-Lindahl et al. (2000) also argue for a role of A β peptide in inhibition of cholinergic neurotransmission, independent of its neurotoxic effect. Given the role of the cholinergic system in facilitating learning and memory, such an action provides a plausible mechanism for POCD, but also carries implications for practice.

RELATIONSHIP OF DIFFERENT ANAESTHETIC AGENTS TO POCD

The studies of POCD pathophysiology thus far have indicated that anaesthetic agents, which promote A β

peptide aggregation, or inhibit cholinergic transmission, are potential mediators of POCD. However, each anaesthetic agent, by virtue of its pharmacology, molecular mass and mode of delivery, is able to perform these actions with different potency. Using nuclear magnetic resonance spectroscopy, Mandal et al. (2009) show that volatile agents of lower molecular mass (isoflurane, desflurane) are more potent promoters of A β peptide oligomerisation, by facilitating interaction of key residues in the peptide. In vivo studies in mice have also demonstrated a calcium-dependent pro-apoptotic effect of isoflurane (Xie et al. 2006). Agents of higher molecular mass (thiopental, diazepam) are unable to interact with these residues, and thus, no oligomerisation is seen.

Likewise, anaesthetic agents differ in their ability to inhibit cholinergic transmission, with ketamine and volatile anaesthetic agents acting as potent antagonists at nicotinic acetylcholine receptors (desflurane at M1, sevoflurane at M1 and M3, and isoflurane at M3 receptor subtypes), and barbiturates acting as potent antagonists at muscarinic acetylcholine receptors. Opioids depress cholinergic transmission, and the action of anticholinesterase drugs (physostigmine, pyridostigmine, neostigmine) is dependent upon their concentration in the cerebrospinal fluid (Fodale et al. 2008). By contrast, clinically used concentrations of propofol show no antagonism of the above receptors, although its action may be potentiated by basal forebrain cholinergic dysfunction (Laalou et al. 2008).

In further support of the increased neurotoxicity of volatile anaesthetic agents, is a prospective randomised study from Liu et al. (2013), which assigned elderly patients undergoing surgery to three different protocols of general anaesthesia: sevoflurane (inhaled), propofol (intravenous) and lidocaine (regional epidural). Those who received inhaled sevoflurane showed an accelerated progression of mild cognitive impairment, according to neuropsychological assessment. Nevertheless, the study was limited by its small sample size, and lack of follow-up cerebrospinal fluid analysis. Interestingly, pre-existing Alzheimer's disease neuropathology may increase vulnerability to the neurotoxic effects of sevoflurane, as found in transgenic studies in mice (Lu et al. 2010).

MOVING FORWARD: IMPLICATIONS FOR CLINICAL PRACTICE

Although estimates of incidence of POCD have been hampered by small sample sizes, varying methods of neuropsychological assessment and lack of consensus on threshold for significant cognitive change (Silverstein et al. 2007), it is indisputable that POCD has long-term impacts on sufferers. The International Study of Postoperative Cognitive Dysfunction (ISPOCD) (Moller et al. 2012) found a reduction in daily activities, earlier retirement, increased dependence of social support and increased mortality associated with POCD. General

anaesthesia in children may also lead to neurocognitive impairment, as a result of damage to the developing brain (McCann et al. 2012).

The studies above indicate an increased propensity for low molecular weight, volatile anaesthetic agents to cause POCD (Liu et al. 2013), which must be borne in mind for patients undergoing hip or open cardiac surgery, in particular (Diegeler et al. 2000). In the prevention of POCD, it is also important to consider patient susceptibility; protective anaesthetic and surgical techniques may be recommended in elderly patients or those who are more likely to have a peri-operative inflammatory stress response. Such techniques may involve intra-operative measurements of anaesthetic depth and cerebral oxygenation, which Ballard et al. (2012) describe decrease incidence of POCD and cognitive decline. Nevertheless, these improved anaesthetic techniques may prove futile in those patients who are already vulnerable to the effects of surgical stress.

CONCLUSION

Progress in molecular biology has allowed improved insight into the neuropathology of post-operative cognitive dysfunction (POCD). Use of anaesthetic agents, and volatile agents in particular, has been associated with increased oligomerisation of A β peptide and inhibition of cholinergic neurotransmission, which gives rise to a clinical picture of impairment of learning and cognition. In the prevention of POCD, patient susceptibility factors and peri-operative factors must be considered.

Acknowledgements: None.

Conflict of interest: None to declare.

References

1. Ballard C, Jones E, Gauge N, Aarsland D, Nilsen OB, Saxby BK, Lowery D, Corbett A, Wesnes K, Katsaiti E, Arden J, Amoako D, Prophet N, Purushothaman B, Green D: Optimised anaesthesia to reduce post operative cognitive decline (POCD) in older patients undergoing elective surgery, a randomised controlled trial. *PLoS One* 7, 2012, e37410.
2. Bekker AY, Weeks EJ: Cognitive function after anaesthesia in the elderly. *Best Pract Res Clin Anaesthesiol* 2003; 117:259-272.
3. Bryson GL & Wyand A: Evidence-based clinical update: general anesthesia and the risk of delirium and postoperative cognitive dysfunction. *Can J Anaesth* 2006; 53:669–677.
4. Dabrowski W, Rzecki Z, Czajkowski M, Pilat J, Wacinski P, Kotlinska E, Sztanke M, Sztanke K, Stazka K, Pasternak K: Volatile anesthetics reduce biochemical markers of brain injury and brain magnesium disorders in patients undergoing coronary artery bypass graft surgery. *J Cardio-thorac Vasc Anesth* 2012; 26:395-402.

5. Diegeler A, Hirsch A, Schneider F, et al.: Neuromonitoring and neurocognitive outcome in off-pump versus conventional coronary bypass operation. *The Annals of Thoracic Surgery* 2000; 69:1162–6.
6. Fodale V, Schifilliti D, Praticò C, Santamaria LB: Remifentanyl and the brain. *Acta Anaesthesiologica Scandinavica* 2008; 52:319–26.
7. Hellstrom-Lindahl E: Modulation of beta-amyloid precursor protein processing and tau phosphorylation by acetylcholine receptors. *European Journal of Pharmacology* 2000; 393:255–63.
8. <http://www.nhsconfed.org/resources/key-statistics-on-the-nhs>
9. Jenkins K, Baker AB: Consent and anaesthetic risk. *Anaesthesia* 2003; 58:962–984.
10. Laalou FZ, De Vasconcelos AP, Oberling P, Jeltsch H, Cassel JC, Pain L: Involvement of the basal cholinergic forebrain in the mediation of general (propofol) anaesthesia. *Anesthesiology* 2008; 108:888–96.
11. Liu Y, Pan N, Ma Y, Zhang S, Guo W, Li H, Zhou J, Liu G, Gao M: Inhaled sevoflurane may promote progression of amnesic mild cognitive impairment: A prospective, randomized parallel-group study. *Am J Med Sci* 2013; 345:355–360.
12. Lu Y, Wu X, Dong Y, Xu Z, Zhang Y, Xie Z: Anesthetic sevoflurane causes neurotoxicity differently in neonatal naïve and Alzheimer disease transgenic mice. *Anesthesiology* 2010; 112:1404–1416.
13. Mandal PK, Fodale V: Isoflurane and desflurane at clinically relevant concentrations induce amyloid beta-peptide oligomerization: an NMR study. *Biochemical and Biophysical Research Communications* 2009; 379:716–20.
14. McCann ME, Soriano SG: General anesthetics in pediatric anaesthesia: influences on the developing brain. *Curr Drug Targets* 2012; 13:944–951.
15. Moller JT, Cluitmans P, Rasmussen LS: Long-term postoperative cognitive dysfunction in the elderly ISPOCD1 study: ISPOCD investigators – International Study of Post-Operative Cognitive Dysfunction. *The Lancet* 1998; 351:857–61.
16. Phillips-Bute B, Mathew JP, Blumenthal JA, Grocott HP, Laskowitz DT, Jones RH, Mark DB, Newman MF: Association of neurocognitive function and quality of life 1 year after coronary artery bypass graft (CABG) surgery. *Psychosom Med* 2006; 68:369–375.
17. Silverstein JH, Timberger M, Reich DL, Uysal S: Central nervous system dysfunction after noncardiac surgery and anesthesia in the elderly. *Anesthesiology* 2007; 106:622–8.
18. Wolman RL, Nussmeier NA, Aggarwal A, Kanchuger MS, Roach GW, Newman MF, Mangano CM, Marschall KE, Ley C, Boisvert DM, Ozanne GM, Herskowitz A, Graham SH, Mangano DT: Cerebral injury after cardiac surgery: identification of a group at extraordinary risk. Multicenter Study of Perioperative Ischemia Research Group (McSPI) and the Ischemia Research Education Foundation (IREF) Investigators. *Stroke* 1999; 30:514–522.
19. Xie Z, Dong Y, Maeda U, et al.: The common inhalation anesthetic isoflurane induces apoptosis and increases amyloid beta protein levels. *Anesthesiology* 2006; 104:988–94.
20. Xie Z, Tanzi RE: Alzheimer's disease and post-operative cognitive dysfunction. *Experimental Gerontology* 2006; 41:346–59.

Correspondence:

Nisha Nesaratnam
School of Clinical Medicine, University of Cambridge
Cambridge, UK
E-mail: nn252@cam.ac.uk